


Interleukin 12 p40 subunit (IL-12p40) is a component of the bioactive cytokines IL-12 and IL-23. Existing in either a monomeric or homodimeric form it is induced in excess of the other subunits of IL-12 and IL-23. Most known for providing negative feedback by competing for the IL-12 receptor it also functions as a chemoattractant for macrophages and is associated with several pathogenic inflammatory responses.

Description	
Source	HEK293, Met1-Ser329, with C-terminal His Tag
UniProt	A0A8U0MDC4
Purity	>99% by SDS-PAGE
Predicted Molecular Mass	39.1 kDa
SDS-PAGE	41.7 kDa
Formulation	Lyophilized from 1X PBS with Trehalose
Preparation and Storage	
Reconstitution	Reconstitute at 25 µg/mL in sterile PBS containing at least 0.1% protein
Shipping	Ambient
Storage	5 years at -80°C; 2 years at -20°C or 4°C Upon reconstitution: 5 years at -80°C; 2 years at -20°C
Expiration	See lot specific CoA
	Ferret IL-12p40 (2µg) on a Coomassie stained 4-12% SDS-PAGE gel, reducing conditions.

For Research Use Only

AMPERSAND BIOSCIENCES, LLC

6459 State Route 30, Lake Clear, NY 12945 | PHONE: (518) 891 0204 | info@ampersandbio.com Rev.1