


Interleukin 6 (IL-6) is a soluble cytokine that can affect inflammation, immune response, and hematopoiesis. IL-6 signals through IL-6R, which can occur as a transmembrane or a soluble receptor that is found in serum. IL-6 is generated immediately following environmental stresses such as infections and tissue injuries. Dysregulation of IL-6, which can lead to over production, may contribute to chronic inflammation and autoimmunity.

Description	
Source	HEK293, Met1-Met217, with C-terminal His Tag
UniProt	P41323
Purity	95% by SDS-PAGE
Predicted Molecular Mass	23 kDa
SDS-PAGE	32.6 kDa
Formulation	Lyophilized from 1X PBS with Trehalose
Preparation and Storage	
Reconstitution	Reconstitute at 10 µg/mL in sterile PBS containing at least 0.1% protein
Shipping	Ambient
Storage	2 years at -80°C; 2 years at -20°C or 4°C Upon reconstitution: 2 years at -80°C; 2 years at -20°C
Expiration	2 years from date of receipt
	Canine IL-6 (2µg) on a Coomassie stained 4-12% SDS-PAGE gel, reducing conditions.

Ampersand Biosciences products may not be sequenced, reverse engineered, or investigated in any way to determine composition without the expressed written consent of Ampersand Biosciences.

For Research Use Only

AMPERSAND BIOSCIENCES, LLC

6459 State Route 30, Lake Clear, NY 12945 | PHONE: (518) 891 0204 | info@ampersandbio.com

Rev 1.0