


Interferon Gamma-Induced Protein 10 (IP-10, CXCL10) is a chemokine that is secreted during an immune response, specifically following IFN gamma stimulation. Following activation by IFN gamma IP-10 initiates Th1-cells and if uncontrolled can lead to a cytokine storm and uncontrolled inflammation.

Description	
Immunogen	HEK293 derived full length recombinant hamster IP-10.
Reactivity	Hamster
Source	Mouse monoclonal IgG ₁
Purification	Protein G purified from hybridoma cell culture supernatant
Applications	ELISA (Capture)
Formulation	Lyophilized from PBS with Trehalose Reconstitute in 100µL ddH ₂ O to 1 mg/mL
Shipping	Ambient
Storage	2 years at -20°C to -80°C as supplied 1 month at 4°C after reconstitution with preservative 1 year at -20°C to -80°C after reconstitution
Expiration	2 years from date of receipt
	<p>Hamster IP-10 Luminex Assay</p> <p>Capture: Hamster IP-10 clone HMIP10-2A7 (Cat #M2065)</p> <p>Detection: Hamster IP-10 polyclonal</p> <p>Antigen: Recombinant Hamster IP-10</p>



For Research Use Only

Ampersand Biosciences products may not be sequenced, reverse engineered, or investigated in any way to determine composition without the expressed written consent of Ampersand Biosciences.